



DATA SHEET

Human Synovial Fluid Membrane Fibroblast Primary Cell Culture - Frozen Vial
--

Catalog number: 36069-03

Description: Frozen Ampule (1.2×10^6 cells) of 1×10^6 viable cells upon thawing, shipped with dry-ice. Also available in T25, T75, T150, and T225 tissue culture flask with plated cells, shipped at room temperature. The Clonal selected Human Synovial Fluid Membrane Fibroblast Primary Cell Culture was derived from Rheumatoid Arthritis. They were maintained in Celprogen's Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Medium and sub-cultured every 24 to 48 hours on Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Extracellular Matrix.

Source: Rheumatoid Arthritis

Donors: All donors from which the Cells were derived were pre-screened; donors tested negative for the usual blood donation infectious disease panel (ABO/RH, Hepatitis B Surface Antigen, HIV1 and 2, Syphilis, hepatitis B Core, Human T Lymphocyte Virus 1 and 2, Hepatitis C Virus, Antibody Screen, Nucleic Amplification Test for HIV 1 HCV, West Nile Virus and Antibodies to Trypanosoma cruzi (the agent of Chagas disease).

Mycoplasma test: Negative-PCR and mycoplasma agar methods

Sterility: Negative for bacteria, yeast, and mold

Storage Conditions: Liquid nitrogen vapor phase for frozen Ampule of Human Synovial Fluid Membrane Fibroblast Primary Cell Culture. For plated cells in tissue culture flask, upon receipt of the cells wipe the flask with 70% ethanol and transfer to sterile tissue culture hood. In the tissue culture hood remove the media from the cells and wash the cells with 1X PBS sterile solution, for 2-3 minutes, remove the 1X PBS solution and then Trypsinize. After Trypsinization of the Cells neutralize the Trypsin with equal volume of Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Media with Serum and collect the Cell suspension in sterile conical centrifuge tube in the tissue culture hood. Centrifuge the cell suspension at 100g for 7 minutes in centrifuge. Plate cells 5×10^5 cells per pre-coated flasks with Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Extracellular Matrix for Expansion in Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Medium.

Positive Markers: Undifferentiated CD 34+, Pro-collagen (collagen type 1), Fibroblast specific protein-1, Prolyl 4-hydroxylase



**Morphology
& Proliferation:**

Mixed population of cells with approximately 95% attached cells and the other 5.0% in suspension, need to change cell culture media every day after 48 hours of initial cell culture or when the media starts changing color to slight yellow from pink. Fast growing cell culture, change media with Celprogen's Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Medium with the appropriate Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Extracellular Matrix. Temperature 37⁰C in 5% CO₂ humidified incubator.

Sub-culturing:

1. Thaw the vial with gentle agitation in a 37⁰C water bath or a dry 37⁰C shaking incubator. For water bath thawing keep the O-ring out of the water.
2. Remove the thawed vial and wipe with 70% ethanol. Then transfer to the tissue culture hood.
3. Transfer the vial contents to a sterile centrifuge tube, and gently add pre-warmed Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Media to the centrifuge tube. Use additional Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Media to rinse the vial and transfer the liquid to the centrifuge tube, repeat this once more to ensure you have all the cells transferred to the 15ml centrifuge tube. Centrifuge the cells at 100g for 7 minutes. Remove the supernatant and re-suspend the cell pellet in 500ul of Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Medium.
4. Add the 500ul of cells to T75 flask, pre-coated with Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Extracellular Matrix, with 10ml of Human Synovial Fluid Membrane Fibroblast Primary Cell Culture Complete Growth Medium.
5. Incubate the cells in the T75 flask at 37⁰C in a 5% CO₂ humidified incubator. Perform 100% Media Change every 24 to 48 hours.
6. Medium renewal every other day or 2-3 days, sub-culturing ratio: 1:2 or 1:3 depending on cell density.
7. Refer to protocols, flow diagrams and videos for more detail. <http://celprogen.com/tech.htm>

Freezing Medium: Available for purchase Cat# M36069-03FM

Trypsin: Available for purchase Cat# T1509-014

IX PBS: Available for Purchase Cat# P1408-013

Storage temperature: Liquid nitrogen vapor phase

Product Orders: Before submitting an order you will be asked to read and accept the terms and conditions of Celprogen's Material Transfer Agreement (MTA).

Permits/Forms: In addition to the MTA mentioned above, other CELPROGEN and/or regulatory permits may be required for the transfer of this CELPROGEN material. Anyone purchasing CELPROGEN material is ultimately responsible for obtaining the permits.

3914 DEL AMO BLVD.
SUITE 901
TORRANCE CA 90503
www.celprogen.com



Phone: 310 542 8822
Fax: 310 542 8028
Email: info@celprogen.com
stemcells@celprogen.com

Notices

& Disclaimers:

CELPROGEN products are intended for laboratory research purposes only. They are not intended for use in Humans. The product, Human Synovial Fluid Membrane Fibroblast Primary Cell Culture – Frozen Vial, is established and manufactured by CELPROGEN Inc., and is for Research Use Only. This product is not for re- sale or may not be transferred to a third party prior to written request and approval by CELPROGEN Inc.