



3914 Del Amo Blvd
Suite 901
Torrance, CA 90503
www.celprogen.com

Phone: 310 542 8822
Fax: 310 542 8028
Email: info@celprogen.com
stemcells@celprogen.com

DATA SHEET

Product name: Lung Endothelial Cell Culture -Plated T150

Catalog number: 11051-05-T150

Description:

Lung Endothelial Cell Culture T150 Plated Cells. Also available in T25, T75, and T225 tissue culture flasks with plated Cells, shipped at room temperature or Frozen vial shipped in Dry-ice. They were maintained in Celprogen's Lung Endothelial Cell Culture Complete Growth Medium and sub-cultured every 24 to 48 hours on Lung Endothelial Cell Culture Culture Extra-cellular Matrix.

Source: Mouse Lung

Mycoplasma test: Negative-PCR and mycoplasma agar methods

Sterility: Negative for bacteria, yeast, and mold

Storage Conditions:

Liquid nitrogen vapor phase for frozen Ampule of Lung Endothelial Cell Culture . For plated cells in tissue culture flask, upon receipt of the cells wipe the flask with 70% ethanol and transfer to sterile tissue culture hood. In the tissue culture hood remove the media of the cells and wash the cells with 1X PBS sterile solution, for 2-3 minutes, remove the PBS solution and then trypsinize. After trypsinization of the Cells neutralize the trypsin with equal volume of Lung Endothelial Cell Culture Complete Growth Media with serum and collect the Cell suspension in sterile conical centrifuge tube in the tissue culture hood. Centrifuge the cell suspension at 100g for 7 minutes in centrifuge. Plate cells 5×10^5 cells per pre-coated flasks with Lung Endothelial Cell Culture Extra-cellular Matrix for Expansion in Lung Endothelial Cell Culture Complete Growth Medium.

Tests Performed:

Microbial: Negative

Incorporation of acetylated Low Density Lipoprotein (LDL): Positive

Positive Markers: Incorporation of acetylated Low Density Lipoprotein (LDL): Positive, CD44, PECAM-1, ICAM-2,

**Morphology
& Proliferation:**

Mixed population of cells with approximately 95% attached cells and the other 5.0% in suspension, need to change cell culture media every day after 48 hours of initial cell culture or when the media starts changing color to slight yellow for pink. Fast growing cell culture. Change media with Celprogen's Lung Endothelial Cell Culture Complete Growth Medium with the appropriate Lung Endothelial Cell Culture



3914 Del Amo Blvd
Suite 901
Torrance, CA 90503
www.celprogen.com

Phone: 310 542-8822
Fax: 310 542-8028
Email: info@celprogen.com
stemcells@celprogen.com

Extra-cellular Matrix. Temperature 37⁰C in 5% CO₂ humidified incubator.

Sub-culturing:

1. Thaw the vial with gentle agitation in a 37⁰C water bath or a dry 37⁰C shaking incubator. For water bath thawing keep the O-ring out of the water.
2. Remove the thawed vial and wipe with 70% ethanol. Then transfer to the tissue culture hood.
3. Transfer the vial contents to a sterile centrifuge tube, and gently add pre-warmed Lung Endothelial Cell Culture Growth Media to the centrifuge tube. Use additional Lung Endothelial Cell Culture Complete Media to rinse the vial and transfer the liquid to the centrifuge tube repeat this once more to ensure you have all the cells transferred to the 15ml centrifuge tube. Centrifuge the cells at 1500 RMP for 5minutes. Remove the supernatant and re-suspend the cell pellet in 500ul of Lung Endothelial Cell Culture Complete Growth Medium.
4. Add the 500ul of cells to T150 flask pre-coated with Lung Endothelial Cell Culture Extra-cellular Matrix with 15ml of Lung Endothelial Cell Culture Complete Growth Medium.
5. Incubate the cells in the T150 flask in a 37⁰C in 5% CO₂ humidified incubator. Perform 100% Media Change every 24 to 48 hours.
6. Medium renewal every other or 2-3 days, sub-culturing ratio: 1:3

Freezing Medium: Available for purchase Cat# M11051-05FM

Trypsin: Available for purchase Cat# T1509-014

IX PBS: Available for Purchase Cat# P1408-013

Storage temperature: Liquid nitrogen vapor phase

Product Orders:

Before submitting an order you will be asked to read and accept the terms and conditions of Celprogen's Material Transfer Agreement (MTA).

Permits/Forms:

In addition to the MTA mentioned above, other CELPROGEN and/or regulatory permits may be required for the transfer of this CELPROGEN material. Anyone purchasing CELPROGEN material is ultimately responsible for obtaining the permits.

Notices

& Disclaimers:

CELPROGEN products are intended for laboratory research purposes only. They are not intended for use in humans. This Product Lung Endothelial Cell Culture established and manufactured by CELPROGEN Inc., and is for Research Use Only. This product is not for re-sale or may not be transferred to third party prior to written request and approval by CELPROGEN Inc.